

# Influenza vaccines for Australians

This fact sheet provides information for immunisation providers on seasonal influenza vaccines that are available in Australia in 2021. It can be used in conjunction with the NCIRS fact sheet [Influenza vaccines – frequently asked questions](#) which provides responses to common questions about influenza viruses and seasonal influenza vaccines.

- Influenza is a common cause of hospitalisation and death in Australia.
- Annual influenza vaccination is recommended for all people aged  $\geq 6$  months.
- Annual influenza vaccination is funded under the National Immunisation Program (NIP) for people aged  $\geq 6$  months who are at increased risk of severe influenza. These include:
  - all Aboriginal and/or Torres Strait Islander people aged  $\geq 6$  months
  - all children aged 6 months to  $< 5$  years
  - all adults aged  $\geq 65$  years
  - people with specified medical conditions (refer to [Table 1](#))
  - pregnant women (during any stage of pregnancy).
- The strains used in seasonal influenza vaccines can change from year to year depending on which viruses are predicted to circulate in each upcoming season.
- Influenza vaccine should not be given on the same day as a COVID-19 vaccine. There should be a 14-day gap between these vaccines.
- A new cell-based influenza vaccine (Flucelvax Quad) is available in Australia. This vaccine is not funded under the NIP.
- For adults aged  $\geq 65$  years, the higher-immunogenicity adjuvanted influenza vaccine (Fluad Quad) is preferentially recommended over standard influenza vaccines.
- Providers are reminded that influenza vaccinations given to people of all ages should be reported to the Australian Immunisation Register (AIR).
- Providers are also reminded to vaccinate pregnant women at any time of the year and any stage of pregnancy. Vaccination of pregnant women provides protection to mothers and their newborn infants.

**The Australian Technical Advisory Group on Immunisation (ATAGI) publishes annual [advice on the use of influenza vaccines in Australia](#).**

## The disease

Influenza or ‘the flu’ is an acute viral illness that mainly affects the respiratory system.

## Causative agent

Influenza is caused by influenza viruses classified as type A, B or C.<sup>1</sup> Only influenza A and B viruses are included in seasonal influenza vaccines as they cause the majority of disease in humans.

Type A influenza viruses are further categorised into subtypes based on two kinds of proteins on their surface: haemagglutinin (H) and neuraminidase (N).<sup>1</sup> Type B influenza viruses are categorised into two lineages: Yamagata and Victoria.

Both influenza A and B can be further broken down into different strains.

The genes for the H and N proteins on the virus surface mutate frequently. This results in constant change to influenza viruses. These minor changes to the H and N proteins of both influenza A and B are referred to as 'antigenic drift' and result in new virus strains. Antibody cross-protection against drifted strains is likely to be reduced. If a major change happens in the H or N protein of influenza A, it is called 'antigenic shift'. Previous immunity is usually not adequate against disease from a 'shifted' strain. This creates the potential for a pandemic.

## Transmission

Influenza is spread easily, mainly through large particle droplets produced by sneezing and coughing.<sup>1,2</sup> Droplets containing the influenza virus also settle onto surfaces, and the virus can then pass from hands to the nose, mouth or eyes. People with influenza can be infectious to others from 24 hours before symptoms start until 1 week after the start of symptoms. In previously healthy individuals, symptoms typically subside within 5–8 days.

People of all ages can get influenza. The percentage of people in the general community affected by flu each year is typically 5–10%, but may be up to 20% in some years. This percentage is higher for children, with 10–40% infected each year.<sup>1-3</sup>

Influenza is more easily spread where large numbers of people gather together.<sup>1</sup> As such, infection rates may be 2–3 times higher in closed populations (e.g. childcare centres, aged care facilities, households).<sup>4,5</sup>

## Clinical features

Influenza symptoms usually have a sudden onset. The most common symptoms are:

- fever
- dry non-productive cough
- nasal congestion
- headache
- sore throat
- constitutional complaints such as myalgia, malaise and fatigue.

The older adults may present with atypical symptoms such as malaise and confusion, and more often develop respiratory complications.

Non-respiratory symptoms such as gastrointestinal complaints and calf muscle pain occur more frequently in children than in adults.<sup>1,3</sup>

Although most influenza infections are symptomatically worse and more severe than other viral upper respiratory tract infections, some may be mild.<sup>1-3</sup> Serious complications from influenza occur in a small proportion of people who are infected.<sup>1-3</sup> Complications include pneumonia, myocarditis and neurological complications, which can lead to hospitalisation and death. People at the highest risk of complications from influenza include those with pre-existing medical conditions. However, previously healthy people can also have severe complications.

## Diagnosis

Laboratory tests are required to confirm an influenza infection. The virus can be detected in a nose or throat swab by rapid antigen-based tests, viral culture or more commonly by molecular

methods, such as polymerase chain reaction (PCR). Serological diagnosis of influenza infection can be made by measuring antibodies in blood samples.

## Treatment

Treatment of influenza, including bed rest, pain relief such as aspirin/paracetamol and fluid intake, generally aims to prevent or minimise symptoms.<sup>1-3</sup> Children and adolescents aged <16 years must not be given aspirin or aspirin-containing medications while sick with influenza because of the increased risk of developing Reye syndrome, a condition that causes swelling in the liver and brain.

Antiviral medications such as oseltamivir or zanamivir, which require a prescription, can help reduce the severity and duration of symptoms of influenza. To be most effective, they need to be given within 48 hours of symptom onset.<sup>1</sup>

## Prevention

Vaccination is the only way to specifically prevent influenza infection and its complications (refer to [Who should be vaccinated](#)).

Practising cough etiquette (such as covering the nose and mouth with a tissue when coughing or sneezing) and washing hands before eating can help reduce the chances of getting and passing on the influenza virus.

People who are sick with influenza should stay home from work, school and social gatherings to prevent close contact with and transmission to other people.<sup>1-3,6</sup>

## Who should be vaccinated

Annual influenza vaccination is recommended for all people aged ≥6 months unless contraindicated (refer to [Contraindications](#)).

There are a number of groups that are at increased risk of influenza and its complications. Annual influenza vaccination is strongly recommended for these groups. For some of these groups, seasonal influenza vaccination is funded under the NIP on the basis of demonstrated cost-effectiveness as a public health intervention.<sup>7</sup> However, annual influenza vaccine should be actively promoted for all people, regardless of eligibility for a funded vaccine.

Influenza vaccination is strongly recommended and funded on the NIP for the following groups:

- all Aboriginal and/or Torres Strait Islander people
- all children aged 6 months to <5 years
- all adults aged ≥65 years
- all people aged ≥6 months with medical conditions **listed in [Table 1](#)** which increase the risk of influenza complications
- pregnant women (during any stage of pregnancy).

Influenza vaccination is strongly recommended but not funded on the NIP for the following groups:

- people with certain medical conditions (in addition to those funded on the NIP):
  - Down syndrome
  - obesity (body mass index ≥30 kg/m<sup>2</sup>)
  - chronic liver disease
- residents and staff (including volunteers) of aged care and long-term residential care facilities
- homeless people

- carers and household contacts of those in high-risk groups
- commercial poultry or pork industry workers
- essential services providers
- travellers

Detailed information on influenza vaccine recommendations is provided in the [Australian Immunisation Handbook](#) (refer also to [Additional resources for primary medical care/vaccination providers](#)).

**Table 1: Medical conditions that are associated with an increased risk of influenza complications and for which individuals are eligible for funded vaccine under the NIP\***

Category	Vaccination strongly recommended for (but not limited to) people with the following clinical conditions
Cardiac disease	Cyanotic congenital heart disease Congestive heart failure Coronary artery disease
Chronic respiratory conditions†	Severe asthma (for which frequent medical consultations or the use of multiple medications is required) Cystic fibrosis Bronchiectasis Suppurative lung disease Chronic obstructive pulmonary disease (COPD) Chronic emphysema
Chronic neurological conditions†	Hereditary and degenerative CNS diseases† (including multiple sclerosis) Seizure disorders Spinal cord injuries Neuromuscular disorders
Immunocompromising conditions‡	Immunocompromised due to disease or treatment (e.g. malignancy, transplantation and/or chronic steroid use) Asplenia or splenic dysfunction HIV infection
Diabetes and other metabolic disorders	Type 1 diabetes Type 2 diabetes Chronic metabolic disorders
Renal disease	Chronic renal failure
Haematological disorders	Haemoglobinopathies
Long-term aspirin therapy in children aged 6 months to 10 years	These children are at increased risk of Reye syndrome following influenza infection

\* Further details are provided in the Australian Immunisation Handbook (refer to [Additional resources for primary medical care/vaccination providers](#)).

† People who have any condition that compromises the management of respiratory secretions or is associated with an increased risk of aspiration should be vaccinated.

‡ People with certain immunocompromising conditions (i.e. haematopoietic stem cell transplant, solid organ transplant) receiving influenza vaccine for the first time post transplant are recommended to receive 2 vaccine doses at least 4 weeks apart (irrespective of age) and 1 dose annually thereafter.

## Contraindications

The only absolute contraindications to influenza vaccines are:

- anaphylaxis after a previous dose of any influenza vaccine
- anaphylaxis due to any vaccine component within influenza vaccine

*Note: Egg allergy is not a contraindication to influenza vaccine. People with egg allergy, including anaphylaxis, can be safely vaccinated with a full dose of influenza vaccine.*

## Vaccines

All influenza vaccines used in Australia are quadrivalent.

The 2021 southern hemisphere egg-based seasonal influenza vaccines contain the following strains:

- A (H1N1) – an A/Victoria/2570/2019 (H1N1)pdm09-like virus
- A (H3N2) – an A/Hong Kong/2671/2019 (H3N2)-like virus
- B (Yamagata lineage) – a B/Phuket/3073/2013-like virus
- B (Victoria lineage) – a B/Washington/02/2019-like virus

The 2021 southern hemisphere cell-based seasonal influenza vaccine contains the following strains:

- A (H1N1) – an A/Wisconsin/588/2019 (H1N1)pdm09-like virus
- A (H3N2) – an A/Hong Kong/45/2019 (H3N2)-like virus
- B (Yamagata lineage) – a B/Phuket/3073/2013-like virus
- B (Victoria lineage) – a B/Washington/02/2019-like virus

Vaccines are registered on the basis of evidence of their effectiveness and safety (refer to [Supplementary information](#)). Multiple registered influenza vaccine products are available each year. The age group(s) in which each vaccine can be used and their NIP availability vary. Refer to [Table 2](#) for the available seasonal influenza vaccines by brand and recommended age.

**Table 2: Seasonal influenza vaccines available for use in Australia in the 2021 influenza season, by brand and recommended age**

(from [ATAGI clinical advice on the administration of seasonal influenza vaccines in 2021](#))

Vaccine Registered age group	Vaxigrip Tetra 0.50 mL (Sanofi)	Fluarix Tetra 0.50 mL (GSK)	FluQuadri 0.50 mL (Sanofi)	Afluria Quad 0.50 mL (Seqirus)	Influvac Tetra 0.50 mL (Mylan)	Flucelvax Quad 0.50 mL (Seqirus)	Fluad Quad 0.50 mL (Seqirus)
6 to 35 months (<3 years)	✓	✓	✓	X	X	X	X
≥3 to <5 years	✓	✓	✓	X	✓	X	X
≥5 to <9 years	✓*	✓*	✓	✓*	✓	X	X
≥9 to <65 years	✓*	✓*	✓	✓*	✓	✓	X
≥65 years	✓	✓	✓	✓	✓	✓	✓†

Ticks indicate age at which a vaccine is registered and available. Shaded boxes represent funding under the NIP.

\* Funding only for Aboriginal and Torres Strait Islander people, pregnant women and people who have certain medical conditions.

† Adjuvanted influenza vaccine preferred over standard influenza vaccines.

More detailed information on seasonal influenza vaccines is provided in the [ATAGI advice for immunisation providers regarding the administration of seasonal influenza vaccines in 2021](#).

## Dosage and administration

The preferred route of administration for influenza vaccines is by intramuscular injection; however, these may also be given by the subcutaneous route. The recommended number of vaccine doses varies by age and immune status of the vaccine recipient. Refer to [Table 3](#) for the recommended doses of seasonal influenza vaccine by age.

Although protection provided by influenza vaccine is generally expected to last for the whole season, optimal protection occurs within the first 3 to 4 months after vaccination.<sup>8,9</sup> While influenza continues to circulate, it is never too late to vaccinate.

All influenza vaccines available in Australia may be co-administered with any other vaccine (refer to [Supplementary information, Safety in infants and children](#)). Detailed information on the administration of influenza vaccines, including co-administration and vaccine interchangeability, is provided in the [Australian Immunisation Handbook](#).

### Table 3: Recommended doses of influenza vaccine by age

(from the current [Influenza disease chapter](#) of the Australian Immunisation Handbook)

Age	Dose	Number of doses needed in 1st year of influenza vaccination	Number of doses needed if person received 1 or more doses of influenza vaccine in a previous season
≥6 months to <9 years	0.5 mL	2 (given 4 weeks apart)	1
≥9 years	0.5 mL	1	1
People of any age who have recently had a haematopoietic stem cell transplant or solid organ transplant	0.5 mL	2 (given 4 weeks apart) in 1st year vaccinated after transplant	2 (given 4 weeks apart) in 1st year vaccinated after transplant then 1 annually

### Co-administration with other vaccines

People can get influenza vaccines at any time before or after, or with, most other vaccines.

Currently, getting an influenza vaccine and a COVID-19 vaccine on the same day is not recommended. A minimum interval of 14 days between an influenza vaccine and a COVID-19 vaccine is preferred. Influenza vaccine can be given before or after any dose of a COVID-19 vaccine as long as a minimum interval of 14 days is observed.

People should get the COVID-19 vaccine first if it is available to them, and then the influenza vaccine. However, if the influenza vaccine is available first, people should get that.

This advice may change as further information becomes available. For more information, refer to [ATAGI advice on relative timing of administering influenza and COVID-19 vaccines in 2021](#).

## Supplementary information

### Epidemiology

Influenza is a seasonal disease in temperate regions. Most cases in Australia occur during the winter months of June through September.<sup>10</sup> In the northern hemisphere, influenza usually occurs between December and April, whereas in the tropics, influenza can occur all year round.

Annual influenza epidemics are most often due to a single virus subtype or lineage. However, the circulating subtypes/lineages can vary year to year and different subtypes/lineages may appear sequentially or simultaneously in the same season.<sup>11</sup>

Influenza is an important cause of illness and death. The number of affected people varies considerably from year to year depending on the characteristics of the circulating virus strains and the level of immunity in the population.

It has long been recognised that the impact of influenza is often greatly under-estimated.<sup>6,12</sup> Between 2006 and 2013 (excluding the 2009 pandemic year), an average of 100 deaths and around 5,100 hospitalisations due to influenza occurred annually in Australia.<sup>13</sup>

In the 2017 influenza season, the highest level of activity since the 2009 pandemic year was recorded. Around 1,100 deaths were reported nationally among notified cases of laboratory-confirmed influenza.<sup>14</sup> However, a mathematical modelling study estimated that influenza is likely to be associated with more than 3,000 deaths and 13,500 hospitalisations each year in Australia, just in people aged >50 years.<sup>15</sup>

There are a number of groups who are at a higher risk of influenza and its complications. These groups experience higher illness and death associated with influenza than the rest of the population:

- Adults aged ≥65 years and children aged <5 years have the highest rates of influenza notifications and hospitalisations.<sup>13</sup>
- Aboriginal and Torres Strait Islander people experience a greater disease burden from influenza than non-Indigenous Australians across all age groups.<sup>13,16</sup>
- People with certain underlying medical conditions such as chronic heart, lung and neuromuscular disease, among others, are also at higher risk of severe influenza complications compared with otherwise healthy individuals.<sup>17</sup>
- Pregnant women are more likely than other women to be hospitalised with influenza.<sup>18</sup> Infants born to mothers who contract influenza during pregnancy are at risk of preterm birth and low birth weight.<sup>19</sup>

## Vaccine effectiveness

The level of protection that influenza vaccine provides against influenza virus varies depending on several factors:

- age of the person
- immunocompetent status of the person
- the level and severity of influenza activity
- how good is the match between the virus strains in the vaccine and those circulating in the community.<sup>6</sup>

A systematic review estimated the efficacy of standard trivalent influenza vaccine to be 59% against laboratory-confirmed influenza in healthy adults aged <65 years; efficacy varied by influenza season.<sup>20</sup> Similar levels of protection have been achieved in young children aged 6 to 59 months, with an estimated vaccine effectiveness of 65% against laboratory-confirmed influenza.<sup>21-23</sup>

Clinical trials of a standard quadrivalent influenza vaccine showed equivalent antibody levels (an accepted surrogate for protection against influenza) to a standard trivalent influenza vaccine for the shared strains in adults and children aged >6 months, and added protection against the additional B strain.<sup>24-27</sup>

The effectiveness of standard influenza vaccines is comparatively lower in older adults, especially in those aged ≥65 years.<sup>28,29</sup> In 2020 an adjuvanted quadrivalent influenza vaccine became available to possibly improve the protection provided through a higher antibody response.<sup>14</sup>

In a large post-licensure study of community-dwelling adults aged ≥65 years, the adjuvanted trivalent influenza vaccine was estimated to be around 25% more effective against hospitalisation for influenza or pneumonia than the standard trivalent influenza vaccine.<sup>30</sup>



In 2021, Flucelvax Quad, a cell-based influenza vaccine, is available for use in Australia. This vaccine is indicated in people aged  $\geq 9$  years. The effectiveness of Flucelvax Quad against laboratory-confirmed influenza is similar to that of standard influenza vaccine.<sup>31,32</sup>

## Vaccine safety

The common symptoms after influenza vaccination can mimic influenza infection, but are due to the vaccine's interaction with the immune system. None of the influenza vaccines available in Australia contain live virus, so they cannot cause influenza.

Less than 15% of people who get influenza vaccine get fever, headache, arthralgia and myalgia. Injection site reactions such as swelling, redness and pain are also common. These side effects may commence within a few hours of vaccination and can last for 1–2 days.<sup>6</sup>

In clinical trials, people who received adjuvanted influenza vaccine had a higher rate of injection site reactions in the week following vaccination than those who received standard influenza vaccine (around 35% versus 18%).<sup>33</sup> Less than 1% of local reactions after both adjuvanted influenza vaccine and standard influenza vaccine were severe.<sup>33</sup>

The number of people who experienced injection site and systemic reactions was similar for both cell-based influenza vaccine and standard influenza vaccine.<sup>34,35</sup> Slightly more injection site reactions were reported in adults aged 18–60 years who received standard influenza vaccine than in those who received cell-based influenza vaccine (29% versus 25%).<sup>35</sup> Less than 1% of injection site and systemic reactions following cell-based influenza vaccine were reported as severe.

Surveillance of influenza vaccine safety through active enhanced surveillance systems such as AusVaxSafety showed that in 2018 across all ages low rates of any adverse event (7.5%) and medical attendance ( $<0.7\%$ ) were reported after vaccination.<sup>36</sup> The fever rate after vaccination in children aged  $<5$  years was less than 1.8%. In 2021, AusVaxSafety will conduct active surveillance in people of all ages. Influenza vaccine safety data will be made available on a weekly basis at [www.ausvaxsafety.org.au](http://www.ausvaxsafety.org.au).

More severe immediate adverse effects such as hives, angioedema or anaphylaxis are rare consequences of influenza vaccination.<sup>6,37,38</sup>

A small increased risk of Guillain-Barré syndrome (GBS) was associated historically with one influenza vaccine in the United States in 1976. But since then, close surveillance has shown that GBS has occurred at a very low rate of less than 1 in 1 million doses of influenza vaccine, if at all.<sup>39,40</sup>

## Safety in infants and children

Surveillance of influenza vaccine safety in young children through AusVaxSafety has shown that influenza vaccine is safe in children aged 6 months to  $<5$  years.

In young children, febrile seizures can occur in susceptible children because of fever of any cause. They occur most often among children aged 12–23 months.<sup>41</sup> Febrile seizures related to influenza vaccination are uncommon: a large study of 69,391 children in the United States calculated a febrile seizure rate of 1.4 per 100,000 in vaccinated children aged 6–23 months.<sup>42</sup>

A slightly higher risk of fever and febrile seizures in children aged 6 months to  $<5$  years (especially those aged 12–24 months) has been reported following the concurrent administration of the inactivated influenza vaccine and the 13-valent pneumococcal conjugate vaccine.<sup>43</sup> This increased risk is small; furthermore, a second more recent study has not demonstrated the same association with febrile seizures and these two vaccines.<sup>44</sup> It is acceptable to administer these vaccines concurrently when both vaccines are indicated.<sup>45</sup>

## Safety in pregnant and lactating women

All influenza vaccines in Australia are inactivated vaccines, which can be safely given to pregnant women at any stage during pregnancy. The rate of adverse events after vaccination in pregnant women is no different from the rate in women who are not pregnant. In addition, studies of mother–baby pairs have shown that receiving the influenza vaccine while pregnant does not increase maternal or fetal complications during pregnancy.<sup>46</sup>

Several high-quality studies<sup>47-51</sup> and an extensive review<sup>52</sup> have shown that influenza vaccination during pregnancy:

- is safe for both the mother and her unborn baby
- protects against preterm birth and low birth weight of the baby
- protects the mother and her newborn in the first few months of life when they are most vulnerable.

Standard influenza vaccines are preferred for use in pregnancy over cell-based influenza vaccines as a larger body of evidence supports the safety of standard influenza vaccines in pregnant women. Although the use of cell-based influenza vaccines in pregnancy has not been assessed, there are no theoretical concerns regarding their safety in pregnant women.

Studies of influenza vaccine given to women who are breastfeeding are limited. However, the available evidence shows that vaccination with influenza vaccine in lactating mothers is safe and can provide protection to the infant.<sup>53</sup>

For further information, refer to the NCIRS fact sheet on [Vaccinations during pregnancy](#).

## Additional resources for primary medical care/vaccination providers

- [NCIRS Influenza vaccines for Australians: Frequently asked questions](#)
- [The Australian Immunisation Handbook](#)
- [Australian Technical Advisory Group on Immunisation \(ATAGI\) advice on seasonal influenza vaccines in 2021](#)
- [ATAGI advice on influenza and COVID-19 vaccines](#)
- [Australian Government Department of Health immunisation website](#)

## References

1. Treanor JJ. Influenza viruses, including avian influenza and swine influenza. In: Mandell GL, Bennett JE, Dolin R, eds. Mandell, Douglas, and Bennett's Principles and practice of infectious diseases. 7th ed ed. Philadelphia, PA: Churchill Livingstone; 2010.
2. Department of Health and Human Services, Centers for Disease Control and Prevention (CDC). Influenza. In: Hamborsky J, Kroger A, Wolfe C, eds. Epidemiology and prevention of vaccine-preventable diseases. 13th ed. Washington, DC: Public Health Foundation; 2015.
3. American Academy of Pediatrics. Influenza. In: Kimberlin DW, Brady MT, Jackson MA, Long SS, eds. Red book: 2015 report of the Committee on Infectious Diseases. 30th ed. Elk Grove Village, IL: American Academy of Pediatrics; 2015.
4. Nicholson KG. Clinical features of influenza. *Seminars in Respiratory Infections* 1992;7:26-37.

5. Hurwitz ES, Haber M, Chang A, et al. Effectiveness of influenza vaccination of day care children in reducing influenza-related morbidity among household contacts. *JAMA* 2000;284:1677-82.
6. Fiore AE, Bridges CB, Katz JM, Cox NJ. Inactivated influenza vaccines. In: Plotkin SA, Orenstein WA, Offit PA, eds. *Vaccines*. 6th ed. Philadelphia, PA: Elsevier Saunders; 2012.
7. Nolan TM. The Australian model of immunization advice and vaccine funding. *Vaccine* 2010;28 Suppl 1:A76-83.
8. Belongia EA, Sundaram ME, McClure DL, et al. Waning vaccine protection against influenza A (H3N2) illness in children and older adults during a single season. *Vaccine* 2015;33:246-51.
9. Sullivan SG, Komadina N, Grant K, et al. Influenza vaccine effectiveness during the 2012 influenza season in Victoria, Australia: influences of waning immunity and vaccine match. *Journal of Medical Virology* 2014;86:1017-25.
10. Chiu C, Dey A, Wang H, et al. Vaccine preventable diseases in Australia, 2005 to 2007. *Communicable Diseases Intelligence* 2010;34 Suppl:ix-S167.
11. Nicholson KG, Wood JM, Zambon M. Influenza. *The Lancet* 2003;362:1733-45.
12. Newall AT, Scuffham PA. Influenza-related disease: the cost to the Australian healthcare system. *Vaccine* 2008;26:6818-23.
13. Li-Kim-Moy J, Yin JK, Patel C, et al. Australian vaccine preventable disease epidemiological review series: Influenza 2006 to 2015. *Communicable Diseases Intelligence* 2016;40:E482-95.
14. Sheridan SL, Patel C, Macartney K, Cheng AC. New enhanced influenza vaccines for older Australians: what promise do they hold? *Medical Journal of Australia* 2018;209:110-2.
15. Newall AT, Wood JG, Macintyre CR. Influenza-related hospitalisation and death in Australians aged 50 years and older. *Vaccine* 2008;26:2135-41.
16. Naidu L, Chiu C, Habig A, et al. Vaccine preventable diseases and vaccination coverage in Aboriginal and Torres Strait Islander people, Australia 2006–2010. *Communicable Diseases Intelligence* 2013;37 Suppl:S1-95.
17. Mertz D, Kim TH, Johnstone J, et al. Populations at risk for severe or complicated influenza illness: systematic review and meta-analysis. *BMJ* 2013;347:f5061.
18. Mertz D, Geraci J, Winkup J, et al. Pregnancy as a risk factor for severe outcomes from influenza virus infection: A systematic review and meta-analysis of observational studies. *Vaccine* 2017;35:521-8.
19. Rasmussen SA, Jamieson DJ, Bresee JS. Pandemic influenza and pregnant women. *Emerging Infectious Diseases* 2008;14:95-100.
20. Osterholm MT, Kelley NS, Sommer A, Belongia EA. Efficacy and effectiveness of influenza vaccines: a systematic review and meta-analysis. *The Lancet Infectious Diseases* 2012;12:36-44.
21. Blyth CC, Jacoby P, Effler PV, et al. Effectiveness of trivalent flu vaccine in healthy young children. *Pediatrics* 2014;133:e1218-25.
22. Heikkinen T, Heinonen S. Effectiveness and safety of influenza vaccination in children: European perspective. *Vaccine* 2011;29:7529-34.

23. Fu C, He Q, Li Z, et al. Seasonal influenza vaccine effectiveness among children, 2010–2012. *Influenza and Other Respiratory Viruses* 2013;7:1168-74.
24. Kieninger D, Sheldon E, Lin WY, et al. Immunogenicity, reactogenicity and safety of an inactivated quadrivalent influenza vaccine candidate versus inactivated trivalent influenza vaccine: a phase III, randomized trial in adults aged ≥18 years. *BMC Infectious Diseases* 2013;13:343.
25. Domachowske JB, Pankow-Culot H, Bautista M, et al. A randomized trial of candidate inactivated quadrivalent influenza vaccine versus trivalent influenza vaccines in children aged 3–17 years. *Journal of Infectious Diseases* 2013;207:1878-87.
26. Pépin S, Donazzolo Y, Jambrecina A, Salamand C, Saville M. Safety and immunogenicity of a quadrivalent inactivated influenza vaccine in adults. *Vaccine* 2013;31:5572-8.
27. Greenberg DP, Robertson CA, Landolfi VA, et al. Safety and immunogenicity of an inactivated quadrivalent influenza vaccine in children 6 months through 8 years of age. *Pediatric Infectious Disease Journal* 2014;33:630-6.
28. Sullivan SG, Chilver MB, Carville KS, et al. Low interim influenza vaccine effectiveness, Australia, 1 May to 24 September 2017. *Euro surveillance : bulletin European sur les maladies transmissibles = European communicable disease bulletin* 2017;22.
29. Fielding JE, Levy A, Chilver MB, et al. Effectiveness of seasonal influenza vaccine in Australia, 2015: an epidemiological, antigenic and phylogenetic assessment. *Vaccine* 2016;34:4905-12.
30. Mannino S, Villa M, Apolone G, et al. Effectiveness of adjuvanted influenza vaccination in elderly subjects in northern Italy. *American Journal of Epidemiology* 2012;176:527-33.
31. Bruxvoort KJ, Luo Y, Ackerson B, et al. Comparison of vaccine effectiveness against influenza hospitalization of cell-based and egg-based influenza vaccines, 2017-2018. *Vaccine* 2019;37:5807-11.
32. DeMarcus L, Shoubaki L, Federinko S. Comparing influenza vaccine effectiveness between cell-derived and egg-derived vaccines, 2017-2018 influenza season. *Vaccine* 2019;37:4015-21.
33. Novartis Vaccines and Diagnostics Inc. FDA advisory committee briefing document: Flud – seasonal adjuvanted trivalent influenza vaccine (aTIV). Vaccines and Related Biological Products Advisory Committee, meeting date: September 15, 2015. 2015. (Accessed 18 March 2019). [http://wayback.archive-it.org/7993/20170405194039/https://www.fda.gov/downloads/AdvisoryCommittees/Committee MeetingMaterials/BloodVaccinesandOtherBiologics/VaccinesandRelatedBiologicalProductsAdvisoryCommittee/UCM461917.pdf](http://wayback.archive-it.org/7993/20170405194039/https://www.fda.gov/downloads/AdvisoryCommittees/Committee%20MeetingMaterials/BloodVaccinesandOtherBiologics/VaccinesandRelatedBiologicalProductsAdvisoryCommittee/UCM461917.pdf)
34. Nolan T, Chotpitayasunondh T, Capeding MR, et al. Safety and tolerability of a cell culture derived trivalent subunit inactivated influenza vaccine administered to healthy children and adolescents: A Phase III, randomized, multicenter, observer-blind study. *Vaccine* 2016;34:230-6.
35. Ambrozaitis A, Groth N, Bugarini R, et al. A novel mammalian cell-culture technique for consistent production of a well-tolerated and immunogenic trivalent subunit influenza vaccine. *Vaccine* 2009;27:6022-9.

36. Pillsbury AJ, Glover C, Jacoby P, et al. Active surveillance of 2017 seasonal influenza vaccine safety: an observational cohort study of individuals aged 6 months and older in Australia. *BMJ Open* 2018;8:e023263.
37. Greenhawt MJ, Li JT, Bernstein DI, et al. Administering influenza vaccine to egg allergic recipients: a focused practice parameter update. *Annals of Allergy, Asthma and Immunology* 2011;106:11-6.
38. Australasian Society of Clinical Immunology and Allergy (ASCIA). Guidelines for medical practitioners: Influenza vaccination of the egg-allergic individual. September 2010. (Accessed 20 March 2018). <https://www.allergy.org.au/hp/papers/vaccination-of-the-egg-allergic-individual>
39. Burwen DR, Ball R, Bryan WW, et al. Evaluation of Guillain-Barré syndrome among recipients of influenza vaccine in 2000 and 2001. *American Journal of Preventive Medicine* 2010;39:296-304.
40. Australian Technical Advisory Group on Immunisation (ATAGI). The Australian Immunisation Handbook, Australian Government Department of Health. Canberra: Australian Government Department of Health; 2019. (Accessed 20 March 2019). <https://immunisationhandbook.health.gov.au/>
41. Australian Government Department of Health, Therapeutic Goods Administration. Seasonal flu vaccine: Investigation into febrile reactions in young children following 2010 seasonal trivalent influenza vaccination. Status report as at 2 July 2010 (updated 24 September 2010). (Accessed 20 March 2019). <http://www.tga.gov.au/alert/seasonal-flu-vaccine-investigation-febrile-reactions-young-children-following-2010-seasonal-trivalent-influenza-vaccination>
42. Hambidge SJ, Glanz JM, France EK, et al. Safety of trivalent inactivated influenza vaccine in children 6 to 23 months old. *JAMA* 2006;296:1990-7.
43. Tse A, Tseng HF, Greene SK, Vellozzi C, Lee GM. Signal identification and evaluation for risk of febrile seizures in children following trivalent inactivated influenza vaccine in the Vaccine Safety Datalink Project, 2010–2011. *Vaccine* 2012;30:2024-31.
44. Kawai AT, Li L, Kulldorff M, et al. Absence of associations between influenza vaccines and increased risks of seizures, Guillain-Barré syndrome, encephalitis, or anaphylaxis in the 2012–2013 season. *Pharmacoepidemiology and Drug Safety* 2014;23:548-53.
45. Kawai AT, Martin D, Kulldorff M, et al. Febrile seizures after 2010–2011 trivalent inactivated influenza vaccine. *Pediatrics* 2015;136:e848-55.
46. Fell DB, Dodds L, MacDonald NE, Allen VM, McNeil S. Influenza vaccination and fetal and neonatal outcomes. *Expert Review of Vaccines* 2013;12:1417-30.
47. Regan AK, Moore HC, de Klerk N, et al. Seasonal trivalent influenza vaccination during pregnancy and the incidence of stillbirth: population-based retrospective cohort study. *Clinical Infectious Diseases* 2016;62:1221-7.
48. Legge A, Dodds L, MacDonald NE, Scott J, McNeil S. Rates and determinants of seasonal influenza vaccination in pregnancy and association with neonatal outcomes. *Canadian Medical Association Journal* 2014;186:E157-64.
49. Zaman K, Roy E, Arifeen SE, et al. Effectiveness of maternal influenza immunization in mothers and infants. *New England Journal of Medicine* 2008;359:1555-64.

50. Benowitz I, Esposito DB, Gracey KD, Shapiro ED, Vázquez M. Influenza vaccine given to pregnant women reduces hospitalization due to influenza in their infants. *Clinical Infectious Diseases* 2010;51:1355-61.
51. Poehling KA, Szilagyi PG, Staat MA, et al. Impact of maternal immunization on influenza hospitalizations in infants. *American Journal of Obstetrics and Gynecology* 2011;204(6 Suppl):S141-8.
52. Giles ML, Krishnaswamy S, Macartney K, Cheng A. The safety of inactivated influenza vaccines in pregnancy for birth outcomes: a systematic review. *Human vaccines & immunotherapeutics* 2018:1-13.
53. Brady RC, Jackson LA, Frey SE, et al. Randomized trial comparing the safety and antibody responses to live attenuated versus inactivated influenza vaccine when administered to breastfeeding women. *Vaccine* 2018;36:4663-71.